

The study of the composition of the visean coal in the eastern part of the east-european platform (Russia federation) in connection with the prospects of their exploitation using well methods

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© SGEM2018. An important source of hydrocarbons is coal which is often associated with oil deposits. In the Volga-Ural oil and gas province (Russian Federation) coal resources in the Visean deposits of the Carboniferous Period exceed 3.5 billion tons. They are located at depths of about 1 km and are inaccessible to mine operations. At the same time, coals can be exploited by well methods as a source of gaseous raw materials. The power gas from coal seams can be used in the development of deposits of high-viscosity oils in adjacent or overlying oil-bearing deposits. For these purposes it is important to know the composition of the organic and inorganic matter of the coals. Visean coals for the most part are metamorphosed to the bituminous coal stage, but at places retain the properties of brown coal. The studies of the coal metamorphism have shown that the reflectance values of vitrinite (R_o) vary in a rather wide range from 0.44 to 0.73%. This parameter depends on the depth of occurrence and the power of heat fluxes in zones of tectonic disturbances. The mineral substance of fossil coals is important to assess the quality of coals and to develop their processing technologies. The main form of the presence of rare elements in coal is mineral phases of pelite dimension, which are difficult to diagnose by traditional methods. In the studied coals, mineral grains of dimension 0.1-50 μm are presented which are effectively studied by scanning electron microscopy. The main minerals in Visean coals are quartz, calcite, feldspar (mainly albite), gypsum, pyrite; kaolinite predominates in the clayey substance; rutile, magnetite, sphalerite are more rare. Coal deposits, formed within a stable platform, are surrounded by carbonate rock strata. They are characterized by the relative homogeneity of the mineral composition and are impoverished by microcomponents.

<http://dx.doi.org/10.5593/sgem2018/1.1/S01.074>

Keywords

Coal, Composition, Exploitation, Organic and inorganic matter, Power gas

References

- [1] Khasanov, R.R., Larochkina, I.A. Prospects for the joint development of oil and coal deposits in previsean depressions of Volga-Ural oil and gas province. *Neftyanoe Khozyaystvo-Oil Industry* (1), 2013, pp. 36-39, (in Russian)

- [2] Kreinin E.V. Nonconventional hydrocarbon sources. New technologies for their development. Moscow: Publisher: Prospekt, 2016, 208 p. (in Russian)
- [3] Walker L. Underground coal gasification: a clean coal technology ready for development, Austral. Coal Rev., pp 19-21, 1999.
- [4] Laouafa, F., Farret, R., Vidal-Gilbert, S., Kazmierczak, J.-B. Overview and modeling of mechanical and thermomechanical impact of underground coal gasification exploitation. Mitigation and Adaptation Strategies for Global Change. 21(4), pp. 547-576, 2016.
- [5] Moore T.A. Coalbed methane: A review, International Journal of Coal Geology, vol. 101, pp 36-81, 2012.
- [6] Khasanov R.R., Varfolomeev M. A., Emel'yanov D.A., and Rakhimzyanov A.I., Investigation of Thermal Effect on Samples of Coals to Determine the Prospect of Their Utilization as Sources of Gaseous Fuels, Chemistry and Technology of Fuels and Oils, 2(606), pp.3-7, 2018.
- [7] Reed, S.J.B., Electron microprobe analysis and scanning electron microscopy in geology, Cambridge: Cambridge Univ., 2nd ed, 192 pp, 2005.
- [8] Khassanov, R.R., Islamov, A.F. and Bogomolov, A.Kh. Rare-earth mineralization in early Carboniferous coals of the Volga-Ural region. Moscow University Geology Bulletin, vol. 69/issue 4, pp. 251-257, 2014.
- [9] Finkelman, R.B., Modes of occurrence of trace elements in coal, Ph.D. Diss., College Park: Dept. Chem., Univ. Mariland, 1980.
- [10] Yudovich Ya.E., Ketris M.P. Inorganic matter of coals. Ekaterinburg: Ural Branch of the Russian Academy of Sciences, 2002, 423 pp. (in Russian)